

**Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

**Listing of Claims:**

1. (Previously Presented) A method of measuring methyl transferase activity of a polypeptide, said method comprising the steps of:
    - a. contacting a polypeptide selected from the group consisting of:
      - i. a polypeptide comprising the amino acid sequence of SEQ ID NO: 51 (*ZNFN3A1*); and
      - ii. a polypeptide that comprises the amino acid sequence having at least about 95% identity to SEQ ID NO: 51 in which both of NHSCDPN (SEQ ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, wherein the polypeptide has a methyltransferase activity for histone H3;with a substrate to be methylated and a cofactor under conditions capable of methylation of the substrate;
  - b. detecting the methylation level of the substrate; and
  - c. measuring the methyl transferase activity by correlating the methylation level of step (b) with the methyl transferase activity.
2. (Original) The method of claim 1, wherein the substrate is a histone or the fragment thereof comprising an at least methylation region.
3. (Original) The method of claim 1, wherein the methylation region is a histone H3 lysine 4.
4. (Previously Presented) The method of claim 1, wherein the cofactor is a S-adenosyl-L-methionine.

5. (Original) The method of claim 1, wherein the condition capable of methylation of the substrate is provided in the existence of heat shock protein 90A (HSP90A).

6. (Original) The method of claim 1, wherein the polypeptide is contacted with the substrate and cofactor in the presence of an enhancing agent for the methylation.

7. (Original) The method of claim 6, wherein the enhancing agent for the methylation is S-adenosyl homocysteine hydrolase (SAHH).

8. (Previously Presented) A method identifying an agent that modulates methyl transferase activity, said method comprising the steps of:

- a. contacting a polypeptide selected from the group consisting of:
  - i. a polypeptide comprising the amino acid sequence of SEQ ID NO: 51; and
  - ii. a polypeptide that comprises the amino acid sequence having at least about 95% identity to SEQ ID NO: 51 in which both of NHSCDPN (SEQ ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, wherein the polypeptide has a methyltransferase activity for histone H3;

with a substrate to be methylated and a cofactor in the presence of a test compound under conditions capable of methylation of the substrate;

- b. detecting the methylation level of the substrate; and
- c. comparing the methylation level to a control level

wherein an increase or decrease in the methylation level compared to control level indicates that the test compound modulates methyl transferase activity.

9. (Previously Presented) A kit for detecting an activity of a test compound to regulate methyl transferase activity, said kit comprising the components of:

- a. a polypeptide selected from the group consisting of:
  - i. a polypeptide comprising the amino acid sequence of SEQ ID NO: 51; and
  - ii. a polypeptide that comprises the amino acid sequence having at least about 95% identity to SEQ ID NO: 51 in which both of NHSCDPN (SEQ

ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, wherein the polypeptide has a methyltransferase activity for histone H3,

- b. a substrate capable of methylation by the polypeptide of (a),
- c. a cofactor for the methylation of the substrate, and
- d. HSP90A.

10. (Original) The kit of claim 9, wherein the substrate is a histone or the fragment thereof comprising an at least methylation region.

11. (Original) The kit of claim 9, wherein said kit further comprises the element of:

- e. S-adenosyl homocysteine hydrolase (SAHH).

12 (Currently Amended) A method of screening for a compound for treating colorectal cancer or hepatocellular carcinoma, said method comprising the steps of:

- a. identifying the compound having an activity to modulate methyl transferase activity by the method of claim 7, and
- b. selecting a compound that decreases the methylation level of the substrate compared to a control level.

13. (Previously Presented) A method of screening for a compound for treating colorectal cancer or hepatocellular carcinoma, said method comprising the steps of:

- a. contacting a polypeptide selected from the group consisting of:
  - i. a polypeptide comprising the amino acid sequence of SEQ ID NO: 51; and
  - ii. a polypeptide that comprises the amino acid sequence having at least about 95% identity to SEQ ID NO: 51 in which both of NHSCDPN (SEQ ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, wherein the polypeptide has a binding activity to a heat shock protein 90A polypeptide (HSP90A);

with a heat shock protein 90A polypeptide (HSP90A) in the presence of a test compound;

- b. detecting binding between the polypeptide and HSP90A;
- c. comparing the binding of the polypeptide and HSP90A in the presence of the test compound with that in the absence of the test compound, and
- d. selecting a test compound which decreases the binding of the polypeptide and HSP90A.

14. (Previously Presented) A kit for screening for a compound for treating colorectal cancer or hepatocellular carcinoma, said kit comprising the components of:

- a. a polypeptide selected from the group consisting of:
  - i. a polypeptide comprising the amino acid sequence of SEQ ID NO: 51; and
  - ii. a polypeptide that comprises the amino acid sequence having at least about 95% identity to SEQ ID NO: 51 in which both of NHSCDPN (SEQ ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, wherein the polypeptide has a binding activity to a heat shock protein 90A polypeptide (HSP90A); and
- b. HSP90A.

15. (Currently Amended) A method of screening for a compound for treating colorectal cancer or hepatocellular carcinoma, said method comprising the steps of:

- a. contacting a polypeptide comprising the amino acid sequence of SEQ ID NO: 51 or a polypeptide ~~fragment comprising a portion of that~~ comprises an amino acid sequence having at least about 95% identity to SEQ ID NO: 51, wherein the polypeptide fragment comprises in which both of NHSCDPN (SEQ ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, and wherein the polypeptide or polypeptide fragment binds to S-adenosyl-L-methionine, with an S-adenosyl-L-methionine in the presence of a test compound;
- b. detecting binding between the polypeptide ~~or polypeptide fragment~~ and S-adenosyl-L-methionine;

- c. comparing the binding of the polypeptide ~~or polypeptide fragment~~ and S-adenosyl-L-methionine in the presence of the test compound with that in the absence of the test compound, and
- d. selecting a test compound which decreases the binding between the polypeptide ~~or polypeptide fragment~~ and S-adenosyl-L-methionine.

16. (Currently Amended) A kit for screening for a compound for treating colorectal cancer or hepatocellular carcinoma, said kit comprising the components of:

- a. a polypeptide comprising the amino acid sequence of SEQ ID NO: 51 or a polypeptide ~~comprising a fragment of that comprises an amino acid sequence having at least about 95% identity to SEQ ID NO: 51, wherein the fragment comprises in which~~ both of NHSCDPN (SEQ ID NO:52) and GEELTICY (SEQ ID NO:53) are conserved, and wherein the polypeptide ~~or polypeptide fragment~~ binds to S-adenosyl-L-methionine; and
- b. S-adenosyl-L-methionine.

17. (Withdrawn) A composition for alleviating a symptom of colorectal cancer or hepatocellular carcinoma, said composition comprising a pharmaceutically effective amount of a compound that decreases ZNFN3A1-mediated methylation and a pharmaceutically acceptable carrier.

18. (Withdrawn) A method for alleviating a symptom of colorectal cancer or hepatocellular carcinoma comprising contacting a colorectal cancer cell or a hepatocellular carcinoma cell with a pharmaceutically effective amount of a compound that decreases ZNFN3A1-mediated methylation.

19. (Withdrawn) A method for alleviating a symptom of colorectal cancer or hepatocellular carcinoma comprising contacting a colorectal cancer cell or a hepatocellular carcinoma cell with a pharmaceutically effective amount of a compound that decreases an interaction between ZNFN3A1 and HSP90A.

20. (Withdrawn) A method for alleviating a symptom of colorectal cancer or hepatocellular carcinoma comprising contacting a colorectal cancer cell or a hepatocellular carcinoma cell with a pharmaceutically effective amount of a compound that decreases an interaction between ZNFN3A1 and S-adenosyl-L-methionine.